

# Complexity, Combinatorial Positivity, and Newton Polytopes

Anshul Adve University of California, Los Angeles, Colleen Robichaux<sup>†</sup> University of Illinois at Urbana-Champaign,  
Alexander Yong University of Illinois at Urbana-Champaign

## INTRODUCTION

The nonvanishing problem asks if a coefficient of a polynomial is nonzero. Many families of polynomials in algebraic combinatorics admit combinatorial counting rules and simultaneously enjoy having saturated Newton polytopes (SNP). Thereby, in amenable cases, nonvanishing is in the complexity class  $NP \cap coNP$  of problems with “good characterizations”. This suggests a new algebraic combinatorics viewpoint on complexity theory.

This paper focuses on the case of Schubert polynomials. These form a basis of all polynomials and appear in the study of cohomology rings of flag manifolds. We give a tableau criterion for nonvanishing, from which we deduce the first polynomial time algorithm. These results are obtained from new characterizations of the Schubertope, a generalization of the permutahedron defined for any subset of the  $n \times n$  grid, together with a theorem of A. Fink, K. Mészáros, and A. St. Dizier (2018), which proved a conjecture of C. Monical, N. Tokcan, and A. Yong (2017).

## DECISION PROBLEMS

A **decision problem** is a problem with a yes or no answer given some input. Some problems have quick algorithms while others seem to require a lengthy search to reach an answer. To better understand these difference, problems are sorted into complexity classes.

Some complexity classes with examples:

- NP: LP ( $\exists x \geq 0, Ax=b?$ )
- coNP: Primes
- P: LP and Primes
- NP-complete: Graph coloring

**Problem 1** When do these coincide?

- $P \stackrel{?}{=} NP$
- $NP \stackrel{?}{=} coNP$
- $NP \cap coNP \stackrel{?}{=} P$




Figure 1: Many believe the equalities in Problem 1 do not hold, giving the diagram above.

## Nonvanishing

In algebraic combinatorics we often study polynomial families:

$$F_\diamond = \sum_{\alpha} c_{\alpha, \diamond} x^\alpha = \sum_{s \in S} wt(s) \in \mathbb{Z}[x_1, \dots, x_n]$$

**Example 2** Below are a few different families:

- With  $\diamond = \lambda$ , use  $F_\lambda = s_\lambda$ , the Schur polynomial, and  $S = SSYT(\lambda)$ . For instance,  $s_{(2,1)} = x_1^2 x_2 + x_1 x_2^2$  since

$$SSYT((2,1)) = \left\{ \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array} \right\}.$$

- With  $\diamond = G$ , use  $F_G = \chi_G$ , Stanley’s chromatic symmetric polynomial, and  $S = \{\text{proper colorings of } G\}$ .
- With  $\diamond = w \in S_\infty$ , use  $F_w = \mathfrak{S}_w$ , but there are many choices for  $S$ .

In this framework, we can discuss the complexity of the **nonvanishing problem**:

**Problem 3** What is the complexity of deciding  $c_{\alpha, \diamond} \neq 0$ , as measured in the input size of  $\alpha$  and  $\diamond$ ?

In our cases of interest,  $c_{\alpha, \diamond} \in \mathbb{Z}_{\geq 0}$  has **combinatorial positivity**, which implies  $\text{nonvanishing}(F_\diamond) \in NP$ .

## NEWTON POLYTOPES

To  $F_\diamond$ , we can associate its **Newton polytope**:

$$\text{Newton}(F_\diamond) = \text{conv}\{\alpha : c_{\alpha, \diamond} \neq 0\} \subseteq \mathbb{R}^n$$

C. Monical-N. Tokcan-A. Yong ’17 defined that  $F_\diamond$  has **saturated Newton polytope (SNP)** if

$$\beta \in \text{Newton}(F_\diamond) \iff c_{\beta, \diamond} \neq 0.$$

**Example 4** Let  $f = x_1 x_2^3 + x_1^3 x_2^2 + x_1 x_2^2 + x_1 x_2$ . Then  $f$  does not have SNP since  $(2,2) \in \text{Newton}(f)$  but  $x_1^2 x_2^2$  does not appear in the monomial expansion of  $f$ .




Figure 2:  $\text{Newton}(f)$  of  $f$  in Example 4

SNP combined with a polynomial-size halfspace description of  $\text{Newton}(F_\diamond)$  implies  $\text{nonvanishing}(F_\diamond) \in coNP$ . Therefore, in many cases  $\text{nonvanishing}(F_\diamond) \in NP \cap coNP$ .

**Example 5** Below is an example and non-example of SNP:

- $s_\lambda$  has SNP and  $\text{nonvanishing}(s_\lambda) \in P$
- $\chi_G$  does not have SNP for  $G$  arbitrary, and  $\text{nonvanishing}(\chi_G) \in NP$ . In fact, for each fixed  $n \geq 3$  it is NP-complete.

If some  $F_\diamond$  with SNP is such that  $\text{Nonvanishing}(F_\diamond)$  is NP – complete, then a polynomial-size halfspace description of  $\text{Newton}(F_\diamond)$  implies

$$coNP \cap NP\text{-complete} \neq \emptyset \implies NP = coNP.$$

## Potential Application

**Conjecture 6 (R. P. Stanley ’95)** If  $G$  is claw-free (i.e., it contains no induced  $K_{1,3}$  subgraph), then  $\chi_G$  is Schur positive.

**Conjecture 7 (C. Monical ’17)** If  $\chi_G$  is Schur positive, then it is SNP.

Combining these gives

**Conjecture 8 (A. Adve-C. Robichaux-A.Yong, ’18)** If  $G$  is claw-free then  $\chi_G$  is SNP.

I. Holyer proved  $n$ -coloring claw-free graphs is NP-complete. Therefore:

An polynomial-size halfspace description proves  $\text{nonvanishing}(\chi_{\text{claw-free } G})$  is coNP. This implies  $NP = coNP$ .

## SCHUBERT POLYNOMIALS

**Schubert polynomials** form a linear basis of all polynomials  $\mathbb{Z}[x_1, x_2, x_3, \dots]$ . They were introduced by A. Lascoux–M.-P. Schützenberger to study the cohomology ring of the flag manifold.

For  $w_0 = n n - 1 \dots 2 1 \in S_n$ ,

$$\mathfrak{S}_{w_0}(x_1, \dots, x_n) := x_1^{n-1} x_2^{n-2} \dots x_{n-1}.$$

Otherwise, for  $w \neq w_0$ , apply Newton’s divided difference operator

$$\partial_i f = \frac{f - f^{s_i}}{x_i - x_{i+1}},$$

recursively using weak Bruhat order to define  $\mathfrak{S}_w$ . To each  $w \in S_\infty$  there is a unique **code**,

$$\text{code}(w) = (c_1, c_2, \dots, c_L) \in \mathbb{Z}_{\geq 0}^L,$$

where  $c_i$  counts the number of boxes in the  $i$ -th row of the Rothe diagram  $D(w)$  of  $w$ .

Let Schubert be  $\text{nonvanishing}(\mathfrak{S}_w)$ . The INPUT is  $\text{code}(w) = (c_1, \dots, c_L)$  with  $c_L > 0$  and  $\alpha \in \mathbb{Z}_{\geq 0}^L$ .

## Theorem 9 (A. Adve-C. Robichaux-A.Yong, ’18)

Schubert  $\in P$ .


For  $w \in S_n$ , let  $\text{Tab}(w, \alpha)$  be the fillings of  $D(w)$  with  $\alpha_i$  many  $k$ ’s, where entries in each column are distinct, and any entry in row  $i$  is  $\leq i$ . We prove Theorem 9 using the following:

## Theorem 10 (A. Adve-C. Robichaux-A.Yong, ’18)

$c_{\alpha, w} \neq 0$  if and only if  $\text{Tab}(D(w), \alpha) \neq \emptyset$ .

In general  $\#\text{Tab}(D(w), \alpha) \geq c_{\alpha, w}$ .

**Example 11** For  $w = 31524$ , the tableaux in the set  $\bigcup_{\alpha} \text{Tab}_{<}(D(w), \alpha)$ :



Hence, for instance,  $c_{(2,1,1), 31524} > 0$  but  $c_{(4), 31524} = 0$ .

Fix  $n \in \mathbb{Z}_{>0}$  and let  $D \subseteq [n]^2$ . We call  $D$  a **diagram** and visualize  $D$  as a subset of an  $n \times n$ .

In 2017, C. Monical-N. Tokcan-A. Yong defined the **Schubertope**  $\mathcal{S}_D$ , a polytope defined for  $D \subseteq [n]^2$ , and conjectured the following:

## Theorem 12 (A. Fink-K. Mészáros-A. St. Dizier, ’17)

$\mathcal{S}_D = \text{Newton}(\mathfrak{S}_w)$ .

Our results give a polynomial time algorithm to check if a lattice point is in  $\mathcal{S}_D$ . This more general result gives a polynomial time algorithm for any polynomial family whose Newton polytopes are Schubertopes.

Additionally, we show that while the nonvanishing problem is easy, the counting problem is hard:

## Theorem 13 (A. Adve-C. Robichaux-A.Yong, ’18)

$c_{\alpha, w}$  is #P-complete.

## Proof Sketch of Theorem 9

- By Theorem 12,  $c_{\alpha, w} \neq 0$  if and only if  $\alpha \in \mathcal{S}_D$ .
- Prove  $\alpha \in \mathcal{S}_D$  if and only if  $\text{Tab}(D, \alpha) \neq \emptyset$ .
- Then introduce a new polytope  $\mathcal{P}(D, \alpha)$  whose integer points biject with  $\text{Tab}(D, \alpha)$ .
- Integer linear programming is hard, but  $\mathcal{P}(D, \alpha)$  is totally unimodular. Now use LP feasibility  $\in P$ .

## CONCLUSION

- We described an algebraic combinatorics paradigm for complexity on theoretical computer science.
- Conversely, complexity gives some new perspectives on algebraic combinatorics.
- We obtain new results about Schubert polynomials and the Schubertope.

## ACKNOWLEDGMENTS

This material is based upon work of CR supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE 1746047. We acknowledge Mathoverflow, S. Kintali’s blog “My brain is open” and R. O’Donnell’s Youtube videos for background. We thank Philipp Hieronymi, Cara Monical, Erik Walsberg and Alexander Woo for helpful conversations.